

1. Masyuk M, Abduelmula A, Morosan-Puopolo G, Ödemis V , Rehim R, Khalida N, Yusuf F, Engele J, Tamamura H, Theiss C, Brand-Saberi B.: Retrograde migration of pectoral girdle muscle precursors depends on CXCR4/SDF-1 signaling. <i>Histochem Cell Biol.</i> 2014;142(5):473-88
2. Lipfert J., Ödemis V. , Wagner DC., Boltze J., Engele J.: CXCR4 and CXCR7 form a functional receptor unit for SDF-1/CXCL12 in primary rodent microglia. <i>Neuropathol Appl Neurobiol.</i> 2013; 39(6): 667-80
3. Lipfert J., Ödemis V. , Engele J.: Grk2 is an Essential Regulator of CXCR7 Signalling in Astrocytes. <i>Cell Mol Neurobiol</i> 2013; 33(1): 111-8
4. Hunger C., Ödemis V. , Engele J.: Expression and function of the SDF-1 chemokine receptors CXCR4 and CXCR7 during limb muscle development and regeneration. <i>Exp. Cell Res.</i> 2012; 318: 2178–2190
5. Glisic D., Lehmann C., Figiel F., Ödemis V. , Lindner R., Engele J.: A novel cross-talk between endothelin and ErbB receptors controlling glutamate uptake into astrocytes. <i>J Neurochem.</i> 2012; 122(4): 844-55
6. Ödemis V. , Lipfert J., Kraft R., Hajek P., Abraham G., Hattermann K., Mentlein R., Engele J.: The presumed atypical chemokine receptor CXCR7 signals through Gi/o proteins in primary rodent astrocytes and human glioma cells. <i>Glia</i> 2012; 60 (3): 372–381
7. Ödemis V. , Boosmann K., Küry P., Engele J.: CXCR7 is an active component of SDF-1 signalling in astrocytes and Schwann cells. <i>J. Cell. Sci.</i> 2010; 123(Pt 7): 1081-8
8. Göttle P., Kremer D., Jander S., Ödemis V. , Engele J., Hartung HP, Küry P.: Activation of CXCR7 receptor promotes oligodendroglial cell maturation. <i>Ann Neurol.</i> 2010 Dec; 68(6): 915-24
9. Kolodziej A., Schulz S., Guyon, A., Wu D.F., Pfeiffer M., Ödemis V. , Holt V., Stumm R.: Tonic Activation of CXC Chemokine Receptor 4 in Immature Granule Cells Supports Neurogenesis in the Adult Dentate Gyrus. <i>J. Neuroscience.</i> 2008; 28 (17): 4488-4500
10. Ödemis V. , Boosmann K., Dieterlen MT, Engele J.: The chemokine SDF-1 controls multiple steps of myogenesis through atypical PKCzeta.. <i>J. Cell. Sci.</i> 2007; 120(Pt 22): 4050-9
11. Ödemis V. , Lamp E., Pezeshki G., Moepps B., Schilling K., Gierschik P., Littman DR, Engele J.: Mice deficient in the chemokine receptor CXCR4 exhibit impaired limb innervation and myogenesis. <i>Mol. Cell. Neurosci.</i> 2005; 30(4): 494-505
12. Ödemis V. , Moepps B., Gierschik P., Engele J.: Interleukin-6 and cAMP induce stromal cell-derived factor-1 chemotaxis in astroglia by up-regulating CXCR4 cell surface expression. Implications for brain inflammation. <i>J. Biol. Chem.</i> 2002; 277(42): 39801-8
13. Ruf S., Ödemis V. , Thasler W., Babl-Weisbarth M., Rose C.: Therapie der Pyoderma fistulans sinifica., <i>Zbl Chir</i> 1998;123:601

14. Evers B., Gerngroß H., **Ödemis V.**: Tourniquetschämie und intramuskulärer Sauerstoffpartialdruck bei Kniegelenksarthroskopien. In: Hartel W. (eds) Klinik und Forschung in der Chirurgie unter dem Aspekt von Effizienz und Ökonomie. Langenbecks Archiv für Chirurgie, vol 1997. Springer, Berlin, Heidelberg

15. Evers B., **Ödemis V.**, Gerngross H.: Intramuscular oxygen partial pressure in patients with chronic exertional compartment syndrome (CECS). Adv. Exp. Med. Biol. 1997; 428: 311-316

16. Evers B., **Ödemis V.**, Gerngross H.: Oxygen partial pressure in the anterior tibial muscle during and after knee surgery with tourniquet control. Adv. Exp. Med. Biol. 1997; 428: 317-325